

DEPARTMENT of ENVIRONMENTAL SERVICES
Water Supply & Pollution Control Division - Biology Bureau

LAKE TROPHIC DATA

MORPHOMETRIC:

Lake: WILSON POND	Lake Area (ha):	4.45
Town: KEENE	Maximum depth (m):	1.8
County: Cheshire	Mean depth (m):	0.9
River Basin: Connecticut	Volume (m ³):	38500
Latitude: 42°56'25" N	Relative depth:	0.8
Longitude: 72°19'15" W	Shore configuration:	1.20
Elevation (ft): 505	Areal water load (m/yr):	195.0
Shore length (m): 900	Flushing rate (yr ⁻¹):	225.0
Watershed area (ha): 1898.5	P retention coeff.:	0.09
% watershed ponded: 0.1	Lake type:	natural w/dam

BIOLOGICAL:

22 January 1992

9 July 1991

DOM. PHYTOPLANKTON (% TOTAL)	#1	NO PHYTOPLANKTON RESULT	SYNEDRA 60%
	#2		
	#3		
PHYTOPLANKTON ABUNDANCE (cells/mL)			840
CHLOROPHYLL-A (µg/L)			5.39
DOM. ZOOPLANKTON (% TOTAL)	#1	NO ZOOPLANKTON RESULTS	SPARSE - NO DOMINANT
	#2		(HORIZONTAL NET TOW -
	#3		NO CELL COUNTS)
ROTIFERS/LITER			
MICROCRUSTACEA/LITER			
ZOOPLANKTON ABUNDANCE (#/L)			
VASCULAR PLANT ABUNDANCE			Very abundant
SECCHI DISK TRANSPARENCY (m)			1.3 Visible on bottom
BOTTOM DISSOLVED OXYGEN (mg/L)		12.9	11.0
BACTERIA (fecal col., #/100 ml) #1			< 10
	#2		< 10
	#3		

SUMMER THERMAL STRATIFICATION:

not stratified

Depth of thermocline (m): None
Hypolimnion volume (m³) : None
Anoxic volume (m³) : None

CHEMICAL:

Lake: WILSON POND

Town: KEENE

	22 January 1992		9 July 1991		
DEPTH (m)	1.0		1.0		
pH (units)	6.5		8.8		
A.N.C. (Alkalinity)	26.0		41.2		
NITRATE NITROGEN	0.75		0.94		
TOTAL KJELDAHL NITROGEN					
TOTAL PHOSPHORUS	0.022		0.019		
CONDUCTIVITY (μ mhos/cm)	195.9		253.8		
APPARENT COLOR (cpu)	29		38		
MAGNESIUM			5.23		
CALCIUM			17.5		
SODIUM			22.4		
POTASSIUM			3.94		
CHLORIDE	23		41		
SULFATE	14		15		
TN : TP					
CALCITE SATURATION INDEX					

All results in mg/L unless indicated otherwise


TROPHIC CLASSIFICATION: 1991

D.O. S.D. PLANT CHL TOTAL CLASS


**	3	6	1	10	Eutro.
----	---	---	---	----	--------

COMMENTS:

1. This pond was not listed in the 1964 report of ponds over 10 acres.
2. Pond level was about 1 to 2 feet below normal.
3. The dominant genera of wholewater plankton were Chroomonas (35%), small pennate diatoms (25%) and Cryptomonas (25%).



[illegible]



[illegible]